



Optimal Laser Treatments for Fungal Nails



ALL LASERS ARE O CREATED EQUAL

A TECHNOLOGY CHANGE IN THE STANDARD OF CARE



A TECHNOLOGY CHANGE IN THE STANDARD OF CARE



Light CAN target by:


- (Photo -Chemical
- Photo -Mechanical and/or
- Photo -Thermal


Processes





Morphology of the Fungi





Outside It's Mostly Chitin

The Anatomy Is

Quite Complex


- Anatomy and Morphology of Fungi
- Mycotic Chromophores
- Thermal relaxation of hyphae and mycelia
- Thermolytic susceptability of fungi
- Photomechanical effects on mycelia
- Spatial distribution of the infection

 in 3D

A Primary Offender



Trichophyton rubrum –T. rubrum





You Don't Need a Laser to Kill Fungus

Many Other Treatments Can Work Just As Well



example:

Boiling Water Works Just Fine

(in the same way as some lasers)



It's NOT Just

Targeting








It's Also...

Anti-Targeting



That Really Matters!

BUT, HOW DO YOU ANTI-TARGET?

what do you need to know?

A Clear Understanding of the "Everything Else"




Understanding

The Environment


- Thermal Properties of Dermal Tissues
- Thermal Properties of the Nail Plate
- Regenerative Properties of the Dermis
- Regenerative Properties of the Nail Matrix
- Thermolytic Susceptability of the Dermis and Matrix
- Optical Properties of the Dermis, Matrix, and Nail Plate

A TYPICAL LONG PULSE LASER

Temperature Profile



>60°C (140°F) for $\approx \frac{1}{2}$ seconds



PROBLEMS WITH OTHER LASER PRODUCTS

Wrong Wavelength

- Can't Penetrate Nail Plate
- Lack of Fungal Chromophores

Wrong Pulse Format

- Long Pulses (ms duration) Lack Selective
 Energy Confinement
- Short Pulses (ns duration) ablate and cause plumes

Wrong Fluence

- Too Much causes pain and side-effects
- Too Little yields little efficacy

Wrong Peak Power

- Ablative or Thermal side-efects
- Reduced efficacy


Wrong Average Power

- Volumetric tissue heating
- Insufficient mycolytic effect

Wrong Spatial Properties

- Pinpoint overtreatment
- Poor coverage of treated areas

Wrong Treatment Protocol



Fungal Survival @ 0.5s



THIS IS WHY

Multiple Treatments Are Necessary, with other Lasers other than Q-Clear™

But May Still Be Ineffective!

Note: At These Durations, Skin Temperatures In Excess of 50°C Can Be Quite **Painful**.

Designed to Anti-Target healthy dermis



and Target what doesn't belong there



including fungus



The Effect of Pulse Format




Figure : Schematic dependence of temperature increases in target and anti-targeted tissue as functions of time for a time-structured laser pulse format given by E(t), the time-dependent laser pulse energy. The dotted lines indicate the temperature levels sufficient to damage the target (T^*_{target}) and anti-target (T^*_{tissue}) , respectively.

Laser Pulse Duration

IS CRITICAL

Laser Temporal Format

IS CRITICAL

OTHERWISE

Unwanted Side Effects

(Pain, Burns, Bleeding, Tissue Spatter)


OR

No Effect



Optimizing The Laser Therapy

- Laser Wavelength
- Laser Temporal Characteristics
- Laser Fluences
- Defining Optimal Treatment Protocols







Clinical RESULTS



Before



After





2



Before

After





After

3




11/8/10 No treatment



4

Before





After



After

5



Before





6

After



Before







SUMMARY OF CINICAL STUDY

SUMMARY OF CLINICAL STUDY

Retrospective study of N=100

- Randomly selected from > 400 treated cases
- Blind graded for nail plate clearance
- Subjects ranged from 3-12 months post Tx

Significant Improvement in > 95% of Subjects

- 0% Pain
- 0% Adverse Side Effects
- 100% Patient Satisfaction



Implication: Essentially all Fungi are killed in a single treatment!

MYCOLOGY

Eradication of nail fungal infections


"On 2 month follow-up 125 patients or 95.42% showed mycological cure (negative microscopy and culture). There was no treatment failure (proximal extension of the lesion during treatment) or other materially adverse side-effect noted.

Clinical cure is associated with the alteration of percentages of disease free nail We find a change of >76% as excellent response, 51-75% as very good response, 26-50 as good response 6-25% as moderate response and >5% as low response to treatment."



Reference: Dr. K. Kalokasidis, et al., Procedings EADV 2013 – to be published, October 2013.

All Subjects 3-12 months Post Tx



All Subjects: Distribution of percentage clearance following Q-Clear™ laser treatment of initially dystrophic nail area (independent of position of dystrophy on the nail plate) for all study subjects at study end. At study end, subjects ranged from 2.8 to 12 months post entry (initial Tx). Note: Due to the metric used, all subjects entering the study (by definition) begin at 0% clearance.



Table 6. Response to Treatment by: gender, age group, onychomycosis severity and fungal type, clinical extent, & location

| All Patients | Patient/Derma tophyte/Nail Group | Excellent response (>75%) | Very good response (50-74) | Good response (25-49) | Moderate response (10-24%) | Low Response (>9%) | No Response (0%) |
|---------------------------------------|--|---------------------------------|-------------------------------------|-----------------------------|----------------------------------|--------------------------|------------------------|
| By Gender | Female | 10 (10.6%) | 44 (46.8%) | 25 (26.6%) | 10 (10.6%) | 0 (0.0%) | 5 (5.3%) |
| | Male | 2 (5.4%) | 9 (24.3%) | 16 (43.2%) | 9 (24.3%) | 0 (0.0%) | 1 (2.7%) |
| | <30 y.o | 3 (15.0%) | 4 (20.0%) | 10 (50.0%) | 2 (10.0%) | 0 (0.0%) | 1 (5.0%) |
| By Age Range | 30-60 | 9 (10.5%) | 37 (43.0%) | 24 (27.9%) | 13 (15.1%) | 0 (0.0%) | 3 (3.5%) |
| | 60> | 0 (0.0%) | 12 (50.0%) | 6 (25.0%) | 4 (16.7%) | 0 (0.0%) | 2 (8.3%) |
| By Severity | Mild | 3 (50.0%) | 2 (33.3%) | 1 (16.7%) | 0 (0.0%) | 0 (0.0%) | 0 (0.0%) |
| | Moderate | 5 (13.5%) | 23 (62.2%) | 9 (24.3%) | 0 (0.0%) | 0 (0.0%) | 0 (0.0%) |
| | Severe | 4 (4.5%) | 28 (31.8%) | 31 (35.2%) | 19 (21.6%) | 0 (0.0%) | 6 (6.8%) |
| | Dermatophytes | 10 (9.3%) | 51 (47.2%) | 38 (35.2%) | 8 (7.4%) | 0 (0.0%) | 1 (0.9%) |
| By Fungal | Candida | 1 (5.3%) | 1 (5.3%) | 3 (15.8%) | 10 (52.6%) | 0 (0.0%) | 4 (21.1%) |
| Туре | Non- dermatophytes | 1 (25.0%) | 1 (25.0%) | 0 (0.0%) | 1 | 0 (0.0%) | 1 (25.0%) |
| | Trich. Rubrum | 9 (9.4%) | 48 (50.0%) | 35 (36.5%) | 4 (4.2%) | 0 (0.0%) | 0 (0.0%) |
| | Distal subungual | 9 (7.3%) | 50 (40.7%) | 40 (32.5%) | 18 (14.6%) | 0 (0.0%) | 6 (4.9%) |
| By Clinical Extent of Infection | Proximal subungual | 0 (0.0%) | 1 (100.0%) | 0 (0.0%) | 0 (0.0%) | 0 (0.0%) | 0 (0.0%) |
| | Superficial white | 1 (50.0%) | 0 (0.0%) | 0 (0.0%) | 1 (50.0%) | 0 (0.0%) | 0 (0.0%) |
| | Dystrophic type | 2 (4.3%) | 14 (29.8%) | 17 (36.2%) | 11 (23.4%) | 0 (0.0%) | 3 (6.4%) |
| | Lateral Edge | 2 (4.3%) | 5 (38.5%) | 5 (38.5%) | 0 (0.0%) | 0 (0.0%) | 1 (7.7%) |
| By Location | Hand | 0 (0.0%) | 1 (9.1%) | 3 (27.3%) | 6 (54.5%) | 0 (0.0%) | 1 (9.1%) |
| - | Feet | 9 (9.9%) | 38 (41.8%) | 27 (29.7%) | 14 (15.4%) | 0 (0.0%) | 3 ()3.3% |







Thank you